



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hence $a+b=49$, $a-b=7$, $a=28$, $b=21$. The geometric verification is evident.

Also solved by B. F. Finkel, A.M., M.Sc., 204 St. Marks Square, Philadelphia, Pa.; J. K. Ellwood, Principal Colfax Sub-District School, Pittsburg, Pa.; A. H. Holmes, Brunswick, Me.; and G. B. M. Zerr, A. M., Ph. D., Parsons, W. Va.; L. E. Newcomb, Los Gatos, California; G. I. Hopkins, Warren High School, Warren, Pa.; J. Scheffer, Hagerstown, Md.

217. Proposed by G. W. DRAKE, Fayetteville, Ark.

If one of the principal axes of a cone which stands on a given base be always parallel to a given right line, the locus of the vertex is an equilateral hyperbola or a right line according as the base is a central conic or a parabola. [Exercise 40, page 94, C. Smith's *Solid Geometry*].

Solution by WILLIAM HOOVER, Ph. D., Professor of Mathematics in the Ohio State University, Athens, O.

Let the given central conic be

$$x^2/a + y^2/b = 1, \quad z=0 \quad \dots\dots\dots (1),$$

and regarded as the central conicoid

$$x^2/a + y^2/b + z^2/c = 1 \quad \dots\dots\dots (2),$$

in which the semi-axis c vanishes.

This solution will employ the principle that the principal axes of a cone, which envelope a given conicoid, are normals to the three confocals which pass through the vertex. The confocals to (2) are given by

$$\frac{x^2}{a+\lambda} + \frac{y^2}{b+\lambda} + \frac{z^2}{c+\lambda} = 1 \quad \dots\dots\dots (3),$$

and so the confocals to (1), by

$$\frac{x^2}{a+\lambda} + \frac{y^2}{b+\lambda} + \frac{z^2}{\lambda} = 1 \quad \dots\dots\dots (4).$$

If (ξ, η, s) be the vertex of the cone, a normal to one of the system (4) is

$$\frac{[a+\lambda][x-\xi]}{\xi} + \frac{[b+\lambda][y-\eta]}{\eta} + \frac{[c+\lambda][z-s]}{s} = 1 \quad \dots\dots\dots (5),$$

and if l, m, n be the direction-cosines of the given line, and ρ an undetermined number, we should have, for the normal to (4) at (η, ξ, s) ,

$$l = \frac{\rho\xi}{a+\lambda}, \quad m = \frac{\rho\eta}{b+\lambda}, \quad n = \frac{\rho s}{\lambda} \quad \dots\dots\dots (6),$$

$$\text{giving } a+\lambda = \frac{\rho\xi}{l}, \quad b+\lambda = \frac{\rho\eta}{m}, \quad \lambda = \frac{\rho s}{n} \quad \dots\dots\dots (7),$$

$$\text{and } l\xi + m\eta + ns = \rho \left(\frac{\xi^2}{a+\lambda} + \frac{\eta^2}{b+\lambda} + \frac{s^2}{\lambda} \right) = \rho \quad \dots\dots\dots (8),$$

since (ξ, η, s) is on (4). From the first two of (7),

$$a-b=\rho[\xi/l-\eta/m] \dots\dots\dots(9),$$

$$\text{and from the first and third of (7), } a=\rho[\xi/l-s/n] \dots\dots\dots(10).$$

$$(8) \text{ and } (9) \text{ give } a-b=[l\xi+m\eta+ns][\xi/l-\eta/m] \dots\dots\dots(11),$$

$$\text{and } (9) \text{ and } (10) \text{ give } [a-b][\xi/l-s/n]=a[\xi/l-\eta/m] \dots\dots\dots(12),$$

showing that the locus of the vertex, when the base is a central conic (1), is the intersection of the hyperboloid (11) by the plane (12), or an equilateral hyperbola.

For the second part, let the base be an indefinitely thin elliptic or hyperbolic paraboloid given by

$$x^2/a+y^2/b=2z \dots\dots\dots(13),$$

and the paraboloids confocal to (13), by

$$\frac{x^2}{a+\lambda} + \frac{y^2}{b+\lambda} = 2z + \lambda \dots\dots\dots(14).$$

The normals to (14) through (ξ, η, s) are given by

$$\frac{[a+\lambda][x-\xi]}{\xi} + \frac{[b+\lambda][y-\eta]}{\eta} = -\frac{z-s}{-1} \dots\dots\dots(15),$$

and we have again

$$l=\frac{\rho\xi}{a+\lambda}, \quad m=\frac{\rho\eta}{b+\lambda}, \quad n=-\rho \dots\dots\dots(16),$$

$$\text{giving } a+\lambda=\rho\xi/l, \quad b+\lambda=\rho\eta/m \dots\dots\dots(17).$$

$$\text{These give } l\xi+m\eta+2ns=\rho\left(\frac{\xi^2}{a+\lambda} + \frac{\eta^2}{b+\lambda} - 2s\right)=\rho\lambda=-n\lambda \dots\dots\dots(18).$$

$$\text{This gives } -\lambda=\frac{l\xi+m\eta+2ns}{n} \dots\dots\dots(19).$$

Adding (19) to the first of (17) and substituting from (16),

$$[l^2-n^2]\xi+lm\eta+2lns-aln=0 \dots\dots\dots(20).$$

Eliminating λ and ρ from (16) and (17),

$$a-b=-n[\xi/l-\eta/m] \dots\dots\dots(21).$$

Therefore in this case the locus of the vertex is a straight line, the intersection of the planes (20) and (21).